

Appl. No. 10/785,985
Amendment dated: November 1, 2005
Reply to OA of: August 1, 2005

Amendments to the Specification:

On page 6, please replace the first full paragraph with the following amended paragraph.

In the oral prophylactic agent of the present invention, the foregoing C-phycocyanin and allophycocyanin are collectively termed “phycocyanin” with concentration ranging from 25 ~ 2000 μ g/mL; the concentration of foregoing spirulina growth factor ranges from 1 ~ 5000 μ g/mL. The most preferable ratio of phycocyanin the total amount of C-phycocyanin and allophycocyanin to spirulina growth factor in aforesaid oral prophylactic agent contained in said water-soluble formula is 2:7.

On page 6, please replace the second full paragraph with the following amended paragraph.

The aforesaid oral prophylactic agent against viral infection can also comprises only water-soluble formula or enteric-coated formula, both of which contain the combination of C-phycocyanin, allophycocyanin, and spirulina growth factor, wherein the percentage of those three ingredients is: 3 ~ 45% C-phycocyanin, 1 ~ 15% allophycocyanin, and 96 ~ 40% spirulina growth factor. The foregoing C-phycocyanin and allophycocyanin are collectively termed “phycocyanin” with concentration ranging between 25 ~ 2000 μ g/mL; the concentration of foregoing spirulina growth factor ranges between 1 ~ 5000 μ g/mL. The most preferable ratio of phycocyanin the total amount of C-phycocyanin and allophycocyanin to spirulina growth factor in aforesaid oral prophylactic agent contained in said water-soluble formula is 2:7.